

P023**Initial experience of laparoscopic right lobectomy after portal vein embolization.**

Hwui-Dong CHO, Ki-Hun KIM*, Dong-Hwan JUNG, Gil-Chun PARK, Young-In YOON, Shin HWANG, Chul-Soo AHN, Deok-Bog MOON, Tae-Yong HA, Gi-Won SONG, Sung-Gyu LEE

Division of Hepatobiliary Surgery and Liver Transplantation, Department of Surgery, Ulsan University and Asan Medical Center, Korea

Introduction : Portal vein embolization is a technique used before hepatic resection to increase the size of remnant liver after surgery. This therapy redirects portal blood to segments of the future liver remnant, resulting in hypertrophy. The purpose of this study was to find out the safety and feasibility of laparoscopic right lobectomy after portal vein embolization.

Methods : All consecutive cases of laparoscopic right lobectomy after portal vein embolization between July 2014 and April 2018 in a tertiary referral hospital were enrolled in this retrospective cohort study. All surgical procedures were performed by one surgeon. There were 14 cases of laparoscopic right lobectomy after portal vein embolization. The group was analysed in terms of donor demographics, preoperative data, postoperative outcomes.

Results : The mean age of the donors was 51.3 ± 6.7 years, the gender ratio for men and women was 18:96. The mean operative time was 402.5 ± 78.8 minutes and mean postoperative hospital stay was 10.1 ± 1.7 days. The number of complications was 6 cases (5.3%) and among them, the Clavien-Dindo classification III or higher complication was 2 (1.8%). There were no mortality cases.

Conclusions : Minimal invasive living donor right hepatectomy was a safe and feasible procedure for donors. It showed an acceptable incidence of complications. The authors suggest that minimal invasive living donor right hepatectomy could be a reasonable operative option for donors in terms of cosmetic effect.

Corresponding Author. : **Ki-Hun KIM** (khkim620@amc.seoul.kr)